



UNIVERSITI PUTRA MALAYSIA

**PATHOGENICITY AND CHARACTERISATION OF NEWCASTLE
DISEASE VIRUS ISOLATED IN IRAN**

MAHDI KIANIZ ADEH

FPV 2002 1

**PATHOGENICITY AND CHARACTERISATION OF NEWCASTLE
DISEASE VIRUS ISOLATED IN IRAN**

By

MAHDI KIANIZADEH

**Thesis Submitted in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy in the Faculty of Veterinary Medicine
Universiti Putra Malaysia**

January 2002



This Thesis is dedicated especially to
My wife and children

Azam Sobhani

and

Hamid

Farzaneh

Hamed

Who have given me strength and courage with their patience
To carry-out this program of research.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Doctor of Philosophy

**PATHOGENICITY AND CHARACTERISATION OF NEWCASTLE
DISEASE VIRUS ISOLATED IN IRAN**

By

MAHDI KIANIZADEH

January 2002

Chairperson : Professor Aini Ideris, Ph.D.

Faculty : Veterinary Medicine

Nine Iranian Iranian Newcastle disease viruses were isolated from different Newcastle disease outbreaks during 1995 to 1999 across Iran. Characterization of the isolates was performed using three standard conventional methods such as mean death time (MDT), intracerebral pathogenicity index (ICPI) and intravenous pathogenicity index (IVPI) in embryonated specific pathogen free (SPF) chicken eggs, day old SPF chicks and 6-week old SPF chickens, respectively. Based on the overall results, all the nine Iranian isolates (MK7, MK12, MK13, MK14, Krd76, Kasra97, GH77, KH2/78 and ES1/99) belong to velogenic pathotype.

Following infection in 6-week old SPF chickens via intra-ocular route, regardless of the isolates, the infected chickens showed depression, ruffled feather and diarrhoea, ending in death. However, there were no obvious respiratory disorders. Meanwhile nervous signs were shown in chickens

inoculated with four isolates (MK7, GH77, MK13 and Kasra97) at the later stage of the infection. Grossly, the initial lesions such as haemorrhage and enlargement were observed in visceral organs including proventriculus, intestine and spleen. No significant changes were observed in the respiratory system. However in some cases, chickens infected with some isolates (MK7, MK12, MK13, Kasra97, KH2/78 and ES1/99) demonstrated limited congestion in the trachea and lungs. Microscopically, the earliest changes were observed in proventriculus, intestine, spleen and liver. Focal and diffuse haemorrhage (proventriculus and intestine), lymphoid depletion (spleen) and mononuclear cell infiltration (liver) were common features. Endotheliosis and malacia were seen in the brain. Based on the MDT results (39.6-56 hours) as well as the presence of lesions in the visceral organs, all the isolates can be grouped as viscerotrop velogenic Newcastle disease virus (VVNDV) pathotype.

The F protein cleavage site gene of all the nine isolates was amplified (242 bp) by RT-PCR and sequenced. Nucleotide and amino acid sequence analysis revealed that the cleavage site of all the isolates consisted of 2 pairs of basic amino acids (Arg) at positions 112,113, 115 and 116 forming the motif ¹¹²RRQRR¹¹⁶, similar to motif in velogenic strains isolated from other countries. However, five amino acids substitutions were observed in isolate MK13 compared to other isolates of which 2 were in F₁ N-terminal, 2 upstream and one downstream of the cleavage site. Two of the amino acids differences in MK13 located at positions 108 and 132 were not observed in published sequence of NDV strains.

Hydropathy profiles were similar for all the isolates. They contained one major hydrophilic area at cleavage site and one major hydrophobic area extended between positions 117 to 142 corresponded to the F₁ N- terminal. This feature of hydropathy profile is similar to that of virulent NDV strains. In phylogenetic analysis, all Iranian isolates are grouped together with other virulent strains in group C. The Russian virulent strain, VOL 95, showed the same amino acid sequence with all the Iranian isolates except MK13 indicating a common origin of the viruses. The importance of these substitution remain to be studied. Finally, based on the results obtained, this study proved that all nine Iranian NDV isolates belonged to the velogenic pathotype.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**KEPATOGENAN DAN PENCIRIAN ISOLAT VIRUS PENYAKIT SAMPAR
AYAM DARIPADA IRAN**

Oleh

MAHDI KIANIZADEH

Januari 2002

Pengerusi : Profesor Aini Ideris, Ph.D.

Fakulti : Perubatan Veterinar

Sembilan virus tempatan penyakit sampar Iran dipencilkan dari beberapa wabak penyakit sampar ayam di Iran dari tahun 1995 hingga 1999. Pencirian setiap isolat dijalankan berdasarkan tiga kaedah konvensional seperti MDT (Mean death time), ICPI (Intracerebral pathogenicity index) dan IVPI di (Intravenous pathogenicity index) dalam telur ayam SPF (specific pathogen free) berembrio, anak ayam SPF berusia satu hari dan ayam SPF berusia 6 minggu. Berdasarkan keputusan keseluruhan yang didapati, kesemua sembilan isolat tempatan tersebut (MK7, MK12, MK13, MK14, Krd76, Kasra97, GH77, KH2/78 dan ES1/99) digolongkan dalam patotip velogenik.

Berikutan daripada jangkitan pada ayam SPF berusia 6 minggu dengan kesemua isolat, melalui cara intra-okular, ayam yang dijangkiti menunjukkan kemurungan, bulu berserabut dan cirit-birit yang membawa kepada kematian. Walau bagaimanapun tiada masalah pernafasan yang ketara. Manakala, tanda kegugupan yang disebabkan oleh beberapa isolat (MK7, GH77, MK13 dan Kasra97) dapat dilihat pada fasa akhir jangkitan. Secara

GH77, MK13 dan Kasra97) dapat dilihat pada fasa akhir jangkitan. Secara kasar, lesi awal seperti hemoraj dan pembesaran organ dapat dilihat pada organ visera termasuk proventrikulus, usus dan limpa. Sistem pernafasan tidak menunjukkan sebarang lesi yang ketara. Walau bagaimanapun, dalam setengah kes ayam yang dijangkiti oleh beberapa isolat (MK7, MK12, MK13, Kasra97, KH2/78 dan ES1/99) menunjukkan sedikit kesebakan pada trakea dan paru-paru. Secara mikroskopik, perubahan awal dapat dilihat pada proventrikulus, usus, limpa dan hati. Hemoraj setempat dan tersebar (proventrikulus dan usus), kemusnahan limfoid (limpa) dan infiltrasi sel mononuklear (hati) merupakan ciri-ciri utama. Endoteliosis dan malasia diperhatikan pada bahagian otak. Berdasarkan keputusan MDT (39.6-56 jam) dan juga kehadiran lesi di organ visera, kesemua isolat tempatan tersebut dikategorikan sebagai VVNDV.

Gen tapak belahan protein F bagi semua isolat diamplifikasi (242bp) melalui RT-PCR dan dijujuk. Analysis jujukan nukleotida dan asid amino mendedahkan bahawa tapak belahan bagi kesemua isolat terdiri daripada 2 pasang asid amino basik (Arg) pada kedudukan 112, 113, 115 dan 116 yang membentuk motif $^{112}\text{RRQRR}^{116}$, menyerupai motif strain velogenik yang dipencil dari negara lain. Walau bagaimanapun, penggantian lima asid amino diperhatikan dalam isolat MK13 berbanding kepada isolat lain yang mana 2 pada terminus-N F1, 2 di hadapan dan 1 dibelakang daripada tapak belahan. Perbezaan dalam 2 asid amino pada MK13 yang terletak pada kedudukan 108 dan 132 tidak diperhatikan dalam jujukan strain NDV yang pernah diterbitkan. Kepentingan penukaran asid amino ini perlu dibuat kajian.

Profil hidropati bagi kesemua isolat adalah serupa. Mereka mempunyai satu kawasan hidrofilik major pada tapak belahan dan satu kawasan hidrofobik major yang menganjur dari kedudukan di antara 117 hingga 142 yang berdekatan dengan terminus-N F1. Ciri profil hidropati ini adalah sama dengan strain NDV virulen. Dalam analisis filogen, kesemua isolat tempatan digolongkan bersama beberapa strain virulent yang lain dalam kumpulan C. Strain virulen Rusia, VOL95, menunjukkan jujukan asid amino yang sama dengan kesemua isolat tempatan kecuali MK13 menandakan persamaan asal-usul bagi virus-virus tersebut. Berdasarkan keputusan yang diperolehi, kajian ini membuktikan bahawa kesemua sembilan isolat NDV Iran tergolong dalam patotip velogenik.

ACKNOWLEDGMENTS

The author gives praises to the almighty Allah for leading him to get this far and making all these possible.

The author is pleased to express his respect, deepest sense and gratitude to Professor Dr. Aini Ideris, Chairperson of advisory committee for her guidance, support, valuable suggestions and encouragement throughout the study period and preparation of this thesis.

The author sincerely expresses his heartfelt gratitude to the members of the advisory committee, Associate Professor Datin Dr. Khatijah Yusoff, Department of Biochemistry and Microbiology, Universiti Putra Malaysia for her guidance and support during this research in her Lab, Dr. Abd Rahman Omar, Biologic Laboratory, Faculty Veterinary Medicine, Universiti Putra Malaysia, for his valuable guidance, constructive suggestions for completion of the research work.

The author wishes to thank co-advisory committee, Professor Dr. M. Shahrabadi, Department of Virology, Iran Medical Science University, Professor Dr. R. Kargar, Department of Virology, Razi Vaccine and Serum Research Institute of Iran, for their advice and assistance during this study.

The author would like thank to Mr. Rajab Beigi, General Director of Education Planning Office and his team in Foreign Students Affairs Department, Iran, for their continue support from the beginning of this study since 1997.

The author expresses his thank to all the staff in Razi Vaccine and Serum Research Institute Iran, for providing facilities during this research.

The author is pleased to express his gratitude and thank to Assistant Professor Dr. Syed Ali Pourbakhsh, Department of Poultry Diseases Diagnosis, Razi Institute, for his guidance and encouragement during this research.

The author extends his gratitudes to Associate Professor Dr. Mohd Hair Bejo, Faculty Veterinary Medicine, Universiti Putra Malaysia for his kind guidance and suggestions.

The author convey his heartfelt thank to his lab mate Omeima Salih, in 1997, for her advice and suggestion in the beginning of this study.

The author wishes to thank Associate Professor Dr. Naderishahab, Department of Plant Molecular Biology, Institute of Rangeland and Forestry in Iran, for his kind guidance and support.

The author expresses his thanks to Dr. Majid Eshaghi, for his kind supports for the phylogenetic analysis.

The author is also grateful to many other, whose names do not appear here, who have helped him in laboratory during the study period.

The author expresses his deepest respect and heartfelt gratitude to his wife for her sacrifices, patience and motivation.

Last but not least, the author owes his greatest gratitude to his beloved mother, father, brothers, sisters, mother and father in-law for their continue support, encouragement and prayers.

I certify that Examination Committee met on 29th November 2001 to conduct the final examination of Mahdi Kianizadeh on his Doctor of Philosophy thesis entitled "Pathogenicity and Characterisation of Newcastle Disease Virus Isolated in Iran" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Siti Suri, Ph.D.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairperson)

Aini Ideris, Ph.D.
Professor.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Khatijah Yusoff, Ph.D.
Associate Professor.
Department of Biochemistry and Microbiology
Universiti Putra Malaysia
(Member)

Abdul Rahman Omar, Ph.D.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Shanshi Shahrabadi, Ph.D.
Professor.
Iran Medical Science University
(Members)

Rohani Kargar Mokhar, Ph.D.
Professor
Raza Vaccine and Serum Research Institute of Iran
(Member)

Nor Muhammad Mahadi, Ph.D.
Professor,
School of Bioscience and Biotechnology
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Independent Examiner)



AINI IDERIS, Ph.D.
Professor/ Dean of Graduate School
Universiti Putra Malaysia

Date: 16 JAN 2002

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy.




AINI IDERIS, Ph.D.
Professor
Dean of Graduate School
Universiti Putra Malaysia

Date: **14** MAR 2007

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



MAHDI KIANIZADEH

16 JAN 2002

TABLE OF CONTENTS

DEDICATION	2
ABSTRACT	3
ABSTRAK	6
ACKNOWLEDGMENTS	9
APPROVAL SHEETS.....	11
DECLARATION FORM.....	13
LIST OF TABLES	18
LIST OF FIGURES	19
LIST OF ABBREVIATIONS	20

CHAPTER

I	INTRODUCTION.....	22
II	LITERATURE REVIEW.....	29
	Newcastle Disease.....	29
	Poultry Industry in Iran.....	31
	Paramyxoviruses.....	33
	Newcastle Disease Virus (NDV).....	36
	Taxonomy.....	36
	Morphology.....	38
	Virus Properties.....	41
	Physical and Chemical Characteristics ...	41
	Epidemiology.....	41
	Host range.....	41
	Vectors.....	42
	Geographical Distribution.....	42
	Speed and Pattern of Spread.....	43
	Vertical Transmission.....	43
	Morbidity and Mortality.....	44
	Pathotypes.....	44
	Pathogenesis.....	45
	Pathology.....	45
	Apoptosis.....	48
	NDV Biologic Properties.....	49
	Agglutinin Activity.....	49
	Cultivation.....	50
	Virulence.....	51
	Pathogenicity Study.....	51
	Clinical signs.....	54
	Diagnosis.....	54
	Molecular Biology of NDV.....	56
	Biologic Properties of Fusion Protein	62
	F Protein Antigenic Epitopes	71
	Polymerase Chain Reaction (PCR).....	73

	Optimisation of the PCR Conditions	74
	Cloning Techniques.....	74
	Ligation.....	76
	Transformation	76
	Screening.....	77
	Sequencing.....	79
	Automatic DNA Sequencing	80
	Modification in Chain Termination Sequencing	80
	Nucleotide Sequencing of NDV	82
	Phylogenetic Analysis	84
	UPGMA Method	84
	Maximum Likelihood Method	84
	Maximum Parsimony	85
III	ISOLATION AND CHARACTERIZATION OF NEWCASTLE DISEASE VIRUS IN IRAN	86
	Introduction.....	86
	Materials and Methods.....	88
	Viruses.....	88
	Isolation Procedure.....	89
	Propagation of Virus	90
	Harvesting of Allantoic Fluid	90
	Clarification of Isolates	90
	Pathogenicity Indices.....	91
	Mean Death Time (MDT).....	91
	Intracerebral Pathogenicity Index	92
	Intravenous Pathogenicity Index	92
	Results	93
	Description of Outbreaks	93
	Isolation	93
	Pathogenicity Indices	95
	MDT Index.....	95
	ICP Index.....	95
	IVP Index.....	95
	Discussion.....	100
IV	PATHOGENESIS OF IRANIAN NDV ISOLATES IN SPF CHICKENS	103
	Introduction.....	103
	Materials and Methods.....	105
	Chickens.....	105
	Virus and Inoculation Procedure.....	
	Sample Collection.....	106
	Sample Preparation for Microscopic Sections	106
	Results.....	106
	Clinical Signs.....	106
	Gross Lesions.....	107

	Microscopic Findings.....	113
	Discussion.....	120
V	MOLECULAR CHARACTERISATION OF IRANIAN NDV FUSION PROTEIN CLEAVAGE SITE	124
	Introduction.....	124
	Materials and Methods.....	127
	Precipitation of Virus.....	127
	Purification of Virus.....	127
	Extraction of RNA	127
	Viral RNA Extraction from Tissues.....	128
	Polymerase Chain Reaction (PCR)	129
	PCR primers.....	129
	Reverse Transcription PCR (RT-PCT).....	129
	PCR Amplification.....	131
	Detection of PCR Product	131
	Purification of PCR Product	131
	Cloning	132
	Ligation	132
	Transformation	135
	Inducing Competency in Cells.....	135
	Antibiotic Resistance Gene Expression.....	136
	Transformant Plating	136
	Screening of Transformed Colonies.....	137
	Analysis of DNA Plasmid Extraction.....	138
	Restriction Enzyme Digestion	138
	Detection of the Segments	138
	Glycerol Stock of Recombinant Plasmids	139
	Sequencing	139
	Kit and Sequencer	139
	Plasmid Extraction	140
	Automatic Sequencer	140
	Phylogenetic Analysis	140
	Results.....	141
	Virus Purification.....	141
	RT-PCR.....	141
	PCR on Tissue Homogenates	141
	Transformant Plating and Selection of Positive Colonies	143
	Restriction Enzyme Digestion	143
	Nucleotide Sequence Analysis.....	145
	Hydropathy Profile	153
	Phylogenetic Analysis.....	154
	Discussion	161
VI	Conclusion.....	170

REFERENCES.....	178
APPENDICES.....	190
Appendix A Pathogenicity Indices of MK7	190
Appendix B Pathogenicity Indices of MK12	191
Appendix C Pathogenicity Indices of MK13	192
Appendix D Pathogenicity Indices of MK14	193
Appendix E Pathogenicity Indices of Krd76	194
Appendix F Pathogenicity Indices of Kasra97	195
Appendix G Pathogenicity Indices of GH77	196
Appendix H Pathogenicity Indices of KH2/78	197
Appendix I Pathogenicity Indices of ES1/99	198
Appendix J Chromas Sequence Result of MK13	199
Appendix K Chromas Sequence Result of Krd76	200
Appendix L Blast search result for KH2/78	201
Appendix M Nucleotide Alignment of the Iranian NDV Isolates	202
Appendix N Comparison of Amino Acid Sequence	203
Appendix O Nucleotide Sequencing Comparison	204
Appendix P Hydrophobicity Scales in Membrane Studies	205
Appendix Q Hydropathy Profile of MK13	206
Appendix R Hydropathy Profile of Krd76	207
Appendix S Hydropathy Profile of La Sota	208
Appendix T DNA Distances Matrices Among 18 NDV Strains	209
Appendix U Distance Matrices of Amino Acids Among 18 NDV Strains	210
VITA	211

LIST OF TABLES

Table	Page
2.1	Records of Newcastle disease outbreaks in Asian countries 30
2.2	Farms related to the broiler line 34
2.3	Farms related to the layer line 34
2.4	Avian Paramyxoviruses 35
2.5	Family of Paramyxoviridae 37
2.6	General properties of the genera Paramyxoviridae 40
2.7	The range of pathogenicity indices of NDV pathotypes 53
2.8	Encoded proteins of NDV 61
2.9	Amino acid sequence at the F ₀ cleavage site 70
3.1	Epizootiological data on NDV outbreaks in Iran 88
3.2	Data on NDV outbreaks in Iran..... 94
3.3	Results of pathogenicity of Iranian isolates 96
3.4	Comparison of ICPI indices among the Iranian isolates 97
3.5	Comparison of IVPI indices among the Iranian isolates 98
3.6	Comparison of pathogenicity indices among the Iranian and some NDV strains 99
4.1	Summarized data on clinical signs in experimental infection 108
4.2	Summarized data on gross lesions 112
4.3	Summarized Data on microscopic changes 119
5.1	Sequence of RT-PCR primers 130
5.2	Primers used for cycle sequencing 139
5.3	Identity range of Iranian NDV isolates in GenBank 146
5.4	Accession numbers assigned to NDV isolate in GenBank 147
5.5	Pathotype of some published NDV strains 152



LIST OF FIGURES

Figure	Page
2.1 Islamic Republic of Iran	32
2.2 Filamentous Form of Negative Stained NDV	39
2.3 Spherical Forms of Negative Stained NDV	39
2.4 Schematic Diagram of Paramyxovirus	59
2.5 Schematic Diagram of F Protein Conformational Changes	64
2.6 Three Major Hydrophobic Area on F Protein	65
2.7 Antigenic Epitopes of F Protein.....	72
4.1 Haemorrhage in Small Intestine	110
4.2 Haemorrhagic Lesions in Proventriculus	110
4.3 Diffuse Haemorrhagic Lesions in Proventriculus	111
4.4 Mononuclear Infiltration in Liver	114
4.5 Normal Intestine Section	115
4.6 Haemorrhagic Intestine Section	115
4.7 Lymphoid Depletion in Spleen	116
4.8 Haemorrhage in Proventriculus	116
4.9 Normal Trachea	117
4.10 Haemorrhage in Trachea	117
4.11 Endotheliosis in Brain	118
4.12 Malacia in Brain	118
5.1 T Vector Genomic Map	134
5.2 Sucrose Gradient of NDV	142
5.3 Detection of Amplified Segment in RT PCR	142
5.4 DNA Plasmid Extraction	144
5.5 DNA Plasmid Digestion	144
5.6 Comparison of Nucleotides Sequence Among the Iranian and some NDV Strains.....	149
5.7 Amino Acid Comparison of Iranian NDV Isolates	151
5.8 Comparison Amino Acid Sequence Among the Iranian and some NDV Strains	151
5.9 Phylogenetic Analysis of Nucleotide Sequences	155
5.10 Phylogenetic Analysis of Amino Acid Sequences	157
5.11 Bootstrap Rooted Tree of Nucleotide Sequences	159
5.12 Bootstrap Rooted Tree of Amino Acid Sequences	160



LIST OF ABBREVIATIONS

A ₂₆₀	Absorption at 260 nm
AI	Avian Influenza
Amp	Ampicillin
Arg	Arginine
bp	Base pairs
CAM	Chorio allantoic membrane
CEC	Council of the economic community
CEF	Chicken embryo fibroblast
CPE	Cytopathic effect
CNS	Central nervous system
Da	Dalton
DEPC	Diethyl pyrocarbonate
DNA	Deoxynucleotide acid
dNTP	Deoxynucleotide triphosphate
ddNTP	Dideoxynucleotide triphosphate
dpi	Days post infection
EDTA	Ethylene diamine tetra acetate
EID	Embryo infection dose
EtOH	Ethanol
ELISA	Enzyme linked immunosorbent assay
F	Fusion protein
GIT	Gastro intestinal tract
HA	Haemagglutinin activity
HI	Haemagglutinin inhibition
HN	Haemagglutinin neuraminidase
hpi	Hours post infection
ICPI	Intra cerebral pathogenicity index
IPTG	Isopropyl-thiogalactopyranoside
IVPI	Intra venous pathogenicity index
Kb	kilo base
KD	kilo dalton
L	Large protein
LB	Luria-bertani
M	Membrane protein
Mabs	Monoclonal antibody
MCS	Multi cloning site
MLD	Minimum lethal dose
mRNA	Messenger RNA
nm	Nanometer
NP	Nucleoprotein
NTE	Sodium chloride-tris-EDTA buffer
OIE	Office International Des Epizooties
ORF	Open reading frame
OD	Optical density

P	Phosphoprotein
Pmol	Picomol
RNA	Ribose nucleic acid
RNase	Ribonuclease
RBCs	Red Blood Cells
SPF	Specific- Pathogen- Free
TAE	Tris-acetate-EDTA buffer
Taq	Thermus aquaticus
V	Valine
VVNDV	Viscerotropic velogenic Newcastle disease virus
X-gal	5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside

CHAPTER I

INTRODUCTION

In the past decades, the systems of poultry husbandry have changed greatly, from small units of birds which were allowed to range freely in fields, to the systems of very large-scale rearing in “broiler house” in which birds are reared intensively or in battery cages for layers. A broiler house may contain as many as 100,000 chickens and there may be a quarter of a million birds on one site. Breeding stocks are also kept intensively, in houses with deep litter.

Keeping such large numbers of birds in close proximity that is less coordinate with physiological system of poultry, adds greatly to the economic loss resulting from a rapid spreading of viral infection of the respiratory system. The control of disease among poultry is further complicated by the present easy transportation of birds and eggs. Among the respiratory diseases, Newcastle disease (ND) has been the major problem in the poultry industry.

Newcastle disease is an infectious ailment of poultry that manifests itself in severe respiratory and nervous conditions. Its causative agent is paramyxovirus, a highly contagious pathogen (Wehmann *et al.*, 1997), known as Newcastle disease virus (NDV). It is an enveloped virus and has a non-segmented negative single stranded RNA genome of approximately 15 kb which codes for six major proteins (Seal *et al.*, 1995).

Outbreaks of ND were first reported in poultry in Java, Indonesia and Newcastle-Upon-Tyne, England in 1926 (Alexander, 1988). From the literature, three panzootics of ND could be recognized. The first appeared in South East Asia in 1926 and spread to most countries of the world. The second panzootic began in the Middle-East in the late 1960s and spread quickly to the other countries by 1973. The third panzootic caused by neurotropic form of ND apparently also started in the Middle-East in the late 1970s, reached Europe by 1981 and spread rapidly worldwide (Alexander, 1988). The disease, currently, has a worldwide distribution with a wide host range in which all orders of birds have been reported to be infected by Newcastle disease virus (NDV). The infectious virus may be ingested or inhaled spreading from one bird to another. This is its major form of transmission and is the basis for mass application of live vaccines via the spray or drinking route (Alexander, 1991).

Isolates of NDV are categorized into three main pathotypes depending on the severity of the disease produced by the isolates in chickens. Lentogenic isolates do not usually cause any disease in adult birds and are considered avirulent. Viruses of intermediate virulence that cause respiratory disease are termed mesogenic while virulent viruses that cause high mortality are termed velogenic. Neurotropic and viscerotropic forms of velogenic viruses have been reported worldwide (Seal *et al.*, 1995).

ND has now been with poultry industry for more than 70 years. Though vaccination for ND has been used successfully from the 1940s (Beard and

Hanson, 1987), its potential to cause great losses in the global poultry industry is probably going to continue as long as this virus can survive and replicate.

In routine diagnosis accessions, NDV isolates are readily identified by haemagglutination-inhibition (HI) with NDV antiserum, and pathogenicity study is often limited to determination of embryo death time (King and Seal, 1998). From the first report of the disease to date, many efforts have been made to develop methods to identify the NDV isolates and to determine their properties. Despite the large number of NDV strains, only minor antigenic differences can be detected with standard serological tests (Marin *et al.*, 1996). Virulence is the main character of NDV that has attracted the attention of researchers in their studies. Among the paramyxoviruses, NDV is unique in that the significant variation in virulence within the same serotype has been described (Waterson *et al.*, 1967).

The role of conventional tests in the determination of virulence of ND viruses is still significant but with some limitations. A source of eggs and chickens, which should preferably be from a specific-pathogen-free flock, is needed. Confirmed diagnosis may be slow, taking several days to isolate the virus and carry out the pathogenicity tests. In addition, the use of animals in this way is becoming increasingly unacceptable with the development of actual or potential alternatives (Aldous and Alexander, 2001). As a result from the progress made in molecular biological techniques especially after the invention of polymerase chain reaction (PCR) by Kary Mullis (1985), attempts have been focused on genetic studies of viral pathogens as well as ND